

CORRECTION Open Access

Lactogenic differentiation of HC11 cells is not accompanied by downregulation of AP-2 transcription factor genes

Richard Jäger*, Leontios Pappas and Hubert Schorle

Following the publication of our article [1], an error in Table 1 was noted. The primer sequences used for PCR of β -casein are incorrectly stated.

The ß-casein primer pair has been incorrectly stated

ß-casein sense: 5'-CCATCCTGCGTCTGGACCTG-3' ß-casein antisense: 5'-GGAATGTTGTGGAGTGG-CAG-3'

The correct ß-casein primer pair that has been used in the study is:

ß-casein sense: 5'-GCCTTGCCAGTCTTGCTAAT-3' ß-casein antisense: 5'-GGAATGTTGTGGAGTGG-CAG-3'

We apologise for any inconvenience this may have caused.

Received: 26 August 2011 Accepted: 9 September 2011 Published: 9 September 2011

Reference

 Jäger R, Pappas L, Schorle H: Lactogenic differentiation of HC11 cells is not accompanied by downregulation of AP-2 transcription factor genes. BMC Res Notes 2008, 23;1:29.

doi:10.1186/1756-0500-4-345

Cite this article as: Jäger *et al.*: Lactogenic differentiation of HC11 cells is not accompanied by downregulation of AP-2 transcription factor genes. *BMC Research Notes* 2011 **4**:345.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



^{*} Correspondence: richard.jaeger@nuigalway.ie Apoptosis Research Centre, School of Natural Sciences, National University of Ireland, Galway, Ireland

